

NASA TECHNICAL
MEMORANDUM

NASA TM X-53104

AUGUST 10, 1964

NASA TM X-53104

N64-31550

FACILITY FORM 804	(ACCESSION NUMBER)	(THRU)
	21	1
	(PAGES)	(CODE)
	TMX-53104	12
	(NASA CR OR TMX OR AD NUMBER)	(CATEGORY)

**AN AUTOMATED MODEL FOR PREDICTING THE
KINETIC TEMPERATURE OF THE AEROSPACE
ENVIRONMENT FROM 100 TO 60,000 KILOMETERS
ABOVE THE SURFACE OF THE EARTH**

by ROBERT E. SMITH
Aero-Astroynamics Laboratory

NASA

*George C. Marshall
Space Flight Center,
Huntsville, Alabama*

OTS PRICE

XEROX	\$	<u>1.00 FS</u>
MICROFILM	\$	<u>0.50 mf</u>

TECHNICAL MEMORANDUM X-53104

AN AUTOMATED MODEL FOR PREDICTING THE KINETIC TEMPERATURE OF THE
AEROSPACE ENVIRONMENT FROM 100 TO 60,000 KILOMETERS ABOVE
THE SURFACE OF THE EARTH

By

Robert E. Smith

George C. Marshall Space Flight Center

Huntsville, Alabama

ABSTRACT

31550

This paper describes the derivation of a series of equations capable of predicting the kinetic temperature of the aerospace environment from 100 to 60,000 kilometers above the surface of the earth. The equations, which are programmed on a GE 225 computer, can be used to predict temperature-height profiles for any time through December 1992.

Author

NASA - GEORGE C. MARSHALL SPACE FLIGHT CENTER

Technical Memorandum X-53104

August 10, 1964

AN AUTOMATED MODEL FOR PREDICTING THE KINETIC TEMPERATURE OF THE
AEROSPACE ENVIRONMENT FROM 100 TO 60,000 KILOMETERS ABOVE
THE SURFACE OF THE EARTH

By

Robert E. Smith

AERO-ASTROPHYSICS OFFICE
SPACE ENVIRONMENT GROUP
AERO-ASTRODYNAMICS LABORATORY
RESEARCH AND DEVELOPMENT OPERATIONS

DEFINITION OF SYMBOLS

<u>Symbol</u>	<u>Definition</u>
a_p	geomagnetic activity index
FT	correction factor for diurnal heating effect
FT1	correction factor for variability of daily mean value of solar flux
FT2	correction factor for variability of the monthly mean value of solar flux
FZ	altitude variation of thermosphere lapse rate
FZ1	altitude variation of correction factor for the diurnal heating effect
S_D	daily mean value of 10.7 centimeter solar radio noise flux ($\times 10^{-22}$ watts per meter 2 per cycle per second)
S_F	monthly mean value of 10.7 centimeter solar radio noise flux ($\times 10^{-22}$ watts per meter 2 per cycle per second)
S_Y	yearly mean value of 10.7 centimeter solar radio noise flux ($\times 10^{-22}$ watts per meter 2 per cycle per second)
SS	annual mean sunspot number
t_L	Local Standard Time in hours and tenths
Z	altitude in kilometers
Z_T	altitude of thermopause in kilometers

TECHNICAL MEMORANDUM X-53104

AN AUTOMATED MODEL FOR PREDICTING THE KINETIC TEMPERATURE OF THE
AEROSPACE ENVIRONMENT FROM 100 TO 60,000 KILOMETERS ABOVE
THE SURFACE OF THE EARTH

SUMMARY

A set of analytical expressions for predicting a time, solar flux, and geomagnetic activity dependent kinetic temperature-height profile of the upper atmosphere above the earth's surface is derived from the data presented by L. G. Jacchia in Smithsonian Institution Astrophysical Observatory Research in Space Science Special Report No. 150, dated 22 April 1964 [1]. Although the data sample is far from adequate and the concept of kinetic temperature in this altitude range is very nebulous, the data that can be provided by this automated predictive model is required to insure the orderly refinement of space vehicle design criteria. Values of kinetic temperature predicted by this model are compared in Figure 1 with the 1959 ARDC Standard and the 1962 U. S. Standard Atmosphere.

I. INTRODUCTION

In a previous paper [2], a model for predicting density-height profiles in the aerospace environment from 200 to 60,000 kilometers above the earth's surface was developed through an analysis of current satellite observations. This is the second model of the series whose final model will be one which predicts the vertical distribution of the molecular weight of the aerospace gas and its temporal and spatial variability. An approach similar to the one employed in Reference 2 was used in this development.

II. DISCUSSION

The first step was the fitting of a curve by the least squares method to the data presented by Jacchia in Reference 1 combined with the kinetic temperature at 100 kilometers quoted in the 1963 Patrick Reference Atmosphere [3] assuming an almost isothermal structure above a thermopause whose height varies with solar activity [1]. Equation (1), below, is a general equation for the temperature-height profile above 100 kilometers above

the earth's surface at 0400 Local Standard Time (LST) when the 10.7 centimeter solar radio noise flux is 67×10^{-22} watts per meter² per cycle per second and the geomagnetic activity index, a_p , is zero.

$$T_{(04, 67, 67, Z)} = (Z - 100) / (0.002022168Z - 0.19844) + 210.00. \quad (1)$$

Equation 2 will predict the height of the thermopause based on the monthly mean value of the 10.7 cm solar radio noise flux assuming a linear variation between Jacchia's heights of 220 kilometers at 70×10^{-22} w/m²-c-s and 400 kilometers at 220×10^{-22} w/m²-c-s [1].

$$Z_T = (6/5) S_F + 135. \quad (2)$$

Equation (3), below, which will predict a kinetic temperature envelope at any altitude, Z, at any time of the day, t_L , for any solar radio noise flux, S_D and S_F , and any geomagnetic activity condition, a_p , incorporates all the corrective factors derived by Jacchia [1].

$$\begin{aligned} T_{(t_L, S_D, S_F, Z, a_p)} &= \left[T_{(04, 67, 67, Z)} + (3.1 \pm 1.9) a_p \right] \left[1 + FT(FZ1) \right] \\ &\quad + FZ \left[FT1 (S_D - 67) + FT2 (S_F - 67) \right]. \end{aligned} \quad (3)$$

The variability of the coefficient of a_p is due to the latitude dependence of the geomagnetic activity effect. The lower value, 1.2, should be used in mid-latitude regions, $30^\circ - 40^\circ$, while the higher value, 5.0, should be used in the auroral zones, $70^\circ - 80^\circ$. If the a_p values are unknown, then zero should be used in the equation and the resulting temperature will be the temperature predicted for a geomagnetically quiet day [4].

In Equation (3), FT is a factor which will adjust the basic temperature-height profile for the diurnal heating effect. The equation for this factor which varies from a minimum of 0.0 at 0400 LST to a maximum of 0.3 at 1400 LST is

$$FT = \sum_{i=0}^{6} a_i \cos i\theta + b_i \sin i\theta \quad (4)$$

where

$$a_0 = 0.1505$$

$$a_1 = 0.14347377$$

$$b_1 = 0.3674753 \times 10^{-1}$$

$$a_2 = 0.8 \times 10^{-2}$$

$$b_2 = -0.11835672 \times 10^{-1}$$

$$a_3 = -0.11666666 \times 10^{-2}$$

$$b_3 = -0.21666666 \times 10^{-2}$$

$$a_4 = 0.15 \times 10^{-2}$$

$$b_4 = -0.13333333 \times 10^{-2}$$

$$a_5 = -0.23071033 \times 10^{-2}$$

$$b_5 = 0.858005 \times 10^{-4}$$

$$a_6 = 0.000.$$

In equation (3), FZ1 is the factor which makes the diurnal heating effect altitude dependent so that the diurnal variation increases from zero at 100 kilometers to 1.3 at and above 120 kilometers.

$$FZ1 = (0.38 - 0.00275Z) (Z - 100.0) \quad (5)$$

for $100 \leq Z = 120$ kilometers and $FZ1 = FZ1$ evaluated at 120 kilometers for $Z > 120$ kilometers.

In equation (3), FZ is a factor which makes the temperature lapse rate in the thermosphere altitude as well as solar activity dependent. The equation for this factor, which varies from 0.00 at 100 kilometers to approximately 0.85 at a 400 kilometer thermopause, is

$$FZ = (Z - 100.0) / (1.1096938Z - 92.828785) \quad (6)$$

for $Z \leq Z_T$ and $FZ = FZ$ evaluated at Z_T for $Z > Z_T$.

FT1, a factor which adjusts the predicted temperature for the daily variation, S_D , in the mean solar radio noise flux, varies from a minimum value of 1.9 at 0400 LST to a maximum value of 2.4 at 1400 LST. If the daily mean value of the 10.7 cm solar flux is unknown, then the value for S_F should be used.

$$FT1 = \sum_{i=0}^{6} c_i \cos i\theta + d_i \sin i\theta \quad (7)$$

where

$$c_0 = 2.15083333$$

$$c_1 = 0.23646679 \quad d_1 = 0.58034166 \times 10^{-1}$$

$$c_2 = 0.14166666 \times 10^{-1} \quad d_2 = -0.24537375 \times 10^{-1}$$

$$c_3 = 0.16666666 \times 10^{-2} \quad d_3 = -0.16666666 \times 10^{-2}$$

$$c_4 = -0.83333333 \times 10^{-3} \quad d_4 = -0.16666666 \times 10^{-2}$$

$$c_5 = -0.31334583 \times 10^{-2} \quad d_5 = 0.29916666 \times 10^{-3}$$

$$c = 0.83333333 \times 10^{-3}.$$

In Equation (3) FT2 is the factor which adjusts the predicted temperature for the variation in the monthly mean solar radio noise flux. The magnitude of this factor varies from a minimum of 4.5 at 0400 LST to a maximum of 6.0 at 1400 LST.

$$FT2 = \sum_{i=0}^{6} e_i \cos i\theta + f_i \sin i\theta \quad (8)$$

where

$$e_0 = 5.26416666$$

$$e_1 = 0.70135016$$

$$f_1 = 0.16649887$$

$$e_2 = 0.50833333 \times 10^{-1}$$

$$f_2 = -0.11833333$$

$$e_3 = -0.83333333 \times 10^{-2}$$

$$f_3 = 0.00000000$$

$$e_4 = -0.91666666 \times 10^{-2}$$

$$f_4 = -0.14433705 \times 10^{-2}$$

$$e_5 = -0.30168333 \times 10^{-2}$$

$$f_5 = 0.13501121 \times 10^{-1}$$

$$e_6 = 0.41666666^{-2}.$$

In equations (4), (7), and (8), $\theta = (t_L - 14.00)\pi/12$ where t_L = Local Standard Time in hours and tenths on a twenty-four hour clock.

The following set of equations was extracted from Reference 2. They will provide the solar activity predictions for use as input data. Equation (9) is used to convert either the known or predicted mean annual sunspot number, SS, to the mean annual value of 10.7 cm solar radio noise flux, S_y . Equations (10a) and (10b) then convert this mean annual value to the mean monthly value, S_M , that would be predicted if the earth's orbit about the sun were circular. Equation (11) corrects this monthly mean value for the eccentricity of the earth's orbit according

to the data presented by Das Gupta [5]. These equations, when combined with a subjective prediction of the annual mean sunspot number from Figure 2, will provide all the data required to compute the predicted temperature-height profile envelopes.

$$S_Y = \sum_{j=0}^8 g_j S^j \quad (9)$$

where

$$g_0 = 67.80036 \quad g_5 = -0.261604186 \times 10^{-8}$$

$$g_1 = 0.414498444 \quad g_6 = -0.159036136 \times 10^{-10}$$

$$g_2 = 0.427501471 \times 10^{-2} \quad g_7 = 0.140461971 \times 10^{-12}$$

$$g_3 = -0.469543503 \times 10^{-4} \quad g_8 = -0.274461310 \times 10^{-15}$$

$$g_4 = 0.660862973 \times 10^{-6}$$

$$S_M = S_Y + \frac{M - 6}{12} (S_{Y+1} - S_Y) \quad \text{when } M \geq 6 \quad (10a)$$

$$S_M = S_Y - \frac{6 - M}{12} (S_Y - S_{Y-1}) \quad \text{when } M < 6 \quad (10b)$$

$$S_F = S_M \sum_{i=0}^6 h_i \cos i\phi + k_i \sin i\phi \quad (11)$$

where

$$h_0 = 1.00$$

$$h_1 = -0.28737186 \times 10^{-1}$$

$$k_1 = -0.13090493 \times 10^{-1}$$

$$h_2 = -0.10333333 \times 10^{-1}$$

$$k_2 = -0.12701714 \times 10^{-1}$$

$$h_3 = 0.150 \times 10^{-2}$$

$$k_3 = -0.350 \times 10^{-2}$$

$$h_4 = -0.50 \times 10^{-3}$$

$$k_4 = -0.20207273 \times 10^{-2}$$

$$h_5 = 0.20601563 \times 10^{-2}$$

$$k_5 = -0.24095063 \times 10^{-2}$$

$$h_6 = 0.28333333 \times 10^{-2}$$

and $\phi = (M - 6) \pi/6$ where M is the month of the year beginning with January = 1, February = 2, etc.

EDP Routine

The following routine is programmed for use on a GE 225 computer. Input data for this program consists of the time of day, t_L , the month of the year, M, the geomagnetic index, a_p , the sunspot number, SS, the daily mean 10.7 cm solar radio noise flux, SD, and the altitude, Z. The output consists of the time of day, TL, the geomagnetic index, AP, the sunspot number, SS, the month of the year, M, the altitude Z, the computed temperature, T, and the 10.7 cm solar radio noise flux, SF.

C MODEL FOR COMPUTING UPPER ATMOSPHERE TEMPERATURE

C R.E. SMITH JOB NUMBER 550120

COMMON SS[30], FM[12], Z[50], SFH[7], SFK[7], GI91,
1F[7], F[7], C[7], D[7], A[7], B[7], SY[30]

P16=.52359878

A=.1505

A[2]=.14347377

A[3]=-8E-2

A[4]=-11666666E-2

A[5]=-15E-2

A[6]=-23071033E-2

A[7]=0.

B=0.

B[2]=-3674753E-1

B[3]=-11835671E-1

B[4]=-21666666E-2

B[5]=-13333333E-2

B[6]=-858005E-4

B[7]=0.

C=2.15083333

C[2]=.23646679

C[3]=.14166666E-1

C[4]=.16666666E-2

C[5]=-83333333E-3

C[6]=-31334583E-2

C[7]=-83333333E-3

D=0.

D[2]=-58034166E-1

D[3]=-24537375E-1

D[4]=-16666666E-2

D[5]=-16666666E-2

D[6]=-29916666E-3

D[7]=0.

E=5.26416666

E[2]=.70135016

E[3]=-50833333E-1

E[4]=-83333333E-2

E[5]=-91666666E-2

E[6]=-30168333E-2

E[7]=-41666666E-2

F=0.

F[2]=-16649887

F[3]=-11833333

F[4]=0.

F[5]=-14433705E-2

F[6]=-13501121E-1

F[7]=0.

G=67.80036

G[2]=-419498444

G[3]=-427501471E-2

G[4]=-469543503E-4

G[5]=-660862973E-6

G[6]=-261604186E-8

G[7]=-159036136E-10

G[8]=-140461971E-12

G[9]=-27446131E-15

C MODEL FOR COMPUTING UPPER ATMOSPHERE TEMPERATURE

```
SFH=1.0
SFH(2)=-.28737186E-1
SFH(3)=-.10333333E-1
SFH(4)=.15E-2
SFH(5)=-.5E-3
SFH(6)=.20601563E-2
SFH(7)=.28333333E-2
SFK=0.
SFK(2)=-.13090493E-1
SFK(3)=-.12701714E-1
SFK(4)=-.35E-2
SFK(5)=-.20207273E-2
SFK(6)=-.24095063E-2
SFK(7)=0.
1 RCD,10P
PRINT 5
5 FORMAT(1H1)
GO TO 15,20,10P
15 RCD,NM,NZ,TL,AP,SD,SS(1),[FM(I),I=1,NM],Z(I),I=1,NZ]
NZ=1
GO TO 30
20 RCD,NM,NS,TL,AP,SD,Z(1),[FM(I),I=1,NM],[SS(I),I=1,NS].
NZ=1
C SY CALCULATION * * * * *
30 DO 35 I=1,NS
SY(I)=G
DO 32 K=2,9
32 SY(I)=SY(I)+G(K)*SS(I)**(K-1)
35 CONTINUE
DO 150 IZ=1,NZ
40 T1=[7/[IZ]=100.1/12.02216E-3*7117]-.19844]+210.
IF([IZ]=120.141,41,42
41 ERR=z(IZ)
GO TO 43
42 ERR=120.
43 CONTINUE
EZ1=[.38,-.00275*ERR]*[ERR=100.1
DO 145 IS=1,NS
DO 140 IM=1,NM
IF([FM(IM)=6,180,50,50
50 IF([IS=NS]70,60,60
60 SM=SY(IS)
GO TO 90
C EQTN 9 A * * * * *
70 SM=SY(IS)+[FM(IM)-6,1/12.*[SY(IS+1)-SY(IS)]
GO TO 90
80 IF([IS=1]60,60,85
85 SM=SY(IS)+[6.-FM(IM)]/12.*[SY(IS)-SY(IS-1)]
90 PHI=[FM(IM)-6,]*PI6
SF=SFH
DO 95 I=2,7
FI=I-1
ANG=FI*PHI
95 SF=SF+SFH(I)*COS(ANG)+SFK(I)*SIN(ANG)
SF=SM*SF
```

C MODEL FOR COMPUTING UPPER ATMOSPHERE TEMPERATURE

```
ZT=6.*SF/5.+135.  
TH=[TL-14.]*PI6/2.  
FT=A  
FT2=E  
FT1=C  
DO 105 I=2,7  
F[I]=I-1  
ANG=F[I]*TH  
SANG=SINF(ANG)  
CANG=CUSF(ANG)  
FT2=FT2+E[I]*CANG+F[I]*SANG  
FT1=FT1+C[I]*CANG+D[I]*SANG  
105 FT=FT+A[I]*CANG+B[I]*SANG  
IF(Z[IZ]-ZT)110,110,120  
110 FZ=(Z[IZ]*100.)/[1.1096938*Z[IZ]-92.828785]  
GO TO 130  
120 FZ=(ZT-100.)/[1.1096938*ZT-92.828785]  
130 IF(SD=1.0)131,131,132  
131 ERR=SF  
GO TO 134  
132 ERR=SD  
134 T=[1.+FT+FZ1)*(T1+1.2*AP)+FZ*(FT1*(ERR=67.)*  
1FT2*(SF=67.))  
PRINT 136,TL,AP,SS(IS),FM(IM1,Z[IZ]),  
1T,SF  
136 FORMAT(///2HTLE17.9,2X2HAP17.9,2X2HSSE17.9,  
12X2HM E17.9//2HZ E17.9,2X2HT E17.9,2X2HSFE17.91  
140 CONTINUE  
145 CONTINUE  
150 CONTINUE  
GO TO 1  
END
```

III. RESULTS

Figure 3 depicts the diurnal variation of the temperature-height profile on a geomagnetically quiet day in the mid-latitudes when the 10.7 cm solar radio noise flux is $100 \times 10^{-22} \text{ w/m}^2\text{-c-s}$.

Figure 4 depicts the diurnal variation of the temperature-height profile on a geomagnetically quiet day in the mid-latitudes when the 10.7 cm solar radio noise flux is $250 \times 10^{-22} \text{ w/m}^2\text{-c-s}$.

Figure 5 depicts the variation with solar activity of the temperature height profile in the mid-latitudes at 0900 LST.

Figure 6 depicts the variation with solar activity of the temperature at mid-latitudes at 0900 LST at constant heights of 200 and 600 kilometers above the earth's surface during the period from July 1964 through December 1992.




FIGURE 1. COMPARISON OF 1959 ARDC, 1962 U.S. STANDARD AND COMPARABLE TEMPERATURES PREDICTED BY THIS MODEL




FIGURE 2. PREDICTED MEAN ANNUAL SUNSPOT NUMBER FOR SOLAR CYCLES 20 AND 21




FIGURE 3. DIURNAL VARIATION IN TEMPERATURE-HEIGHT PROFILE AT $96 \times 10^{-22} \text{W/m}^2\text{-c-s}$
UNITS OF 10.7 CM SOLAR RADIO NOISE FLUX




FIGURE 4. DIURNAL VARIATION IN TEMPERATURE-HEIGHT PROFILE AT $237 \times 10^{-22} \text{W/m}^2\text{-c-s}$
UNITS OF 10.7 CM SOLAR RADIO NOISE FLUX




FIGURE 5. VARIATION IN TEMPERATURE-HEIGHT PROFILE AT 0900 LST WITH SOLAR RADIO NOISE FLUX




FIGURE 6. VARIATION OF TEMPERATURE AT 200 AND 600 KILOMETERS DURING SOLAR CYCLES 20 AND 21

REFERENCES

1. Jacchia, L. G., "The Temperature Above the Thermopause," Smithsonian Institution Astrophysical Observatory Research in Space Science Special Report Number 150, April 22, 1964.
2. Smith, R. E., "An Automated Model for Predicting Aerospace Density Between 200 and 60,000 Kilometers Above the Surface of the Earth," NASA TM X-53062, June 10, 1964, Marshall Space Flight Center, Huntsville, Alabama.
3. Smith, O. E., "Availability of Computer Subroutine for the 1963 Patrick Reference Atmosphere," MSFC Office Memo R-AERO-Y-12-63, December 9, 1963.
4. Jacchia, L. G. and J. Slowey, "Atmospheric Heating in the Auroral Zones: A Preliminary Analysis of the Atmospheric Drag of the Injun 3 Satellite," J. of Geophysical Research 69(5): 905-910, March 1, 1964.
5. Das Gupta, M. K. and D. Basu, "Effect of the Earth's Orbital Eccentricity on Incident Solar Flux at 10.7 cm," J. of the Atmospheric and Terrestrial Physics 26: 135-137, 1964: Pergamon Press Ltd. Printed in Northern Ireland.

August 10, 1964

APPROVAL

NASA TM X-53104


AN AUTOMATED MODEL FOR PREDICTING THE KINETIC TEMPERATURE OF THE
AEROSPACE ENVIRONMENT FROM 100 TO 60,000 KILOMETERS ABOVE
THE SURFACE OF THE EARTH

By

Robert E. Smith


The information in this report has been reviewed for security classification. Review of any information concerning Department of Defense or Atomic Energy Commission programs has been made by the MSFC Security Classification Officer. This report, in its entirety, has been determined to be unclassified.

This document has also been reviewed and approved for technical accuracy.



W. W. Vaughan

Chief, Aero-Astrophysics Office



E. D. Geissler

Director, Aero-Astrodynamic Laboratory

DISTRIBUTION

DEP DIR

Dr. E. Rees

HME-P

R-DIR

Mr. Weidner

R-P&VE

Mr. Cline
Mr. Hellebrand
Dr. Krause
Mr. Goerner (2)

R-SA

Dr. Kuettner

R-COMP

Dr. Hoelzer (2)
Miss Sylvia Bowers

R-FP

Dr. Koelle
Dr. H. O. Ruppe (2)

R-RP

Dr. Stuhlinger (2)
Dr. Johnson
Dr. Shelton

R-AERO

Dr. Geissler
Mr. Jean
Dr. Speer
Mr. Kurtz (3)
Mrs. A. McNair
Mr. J. Lindberg
Dr. Hoelker
Mr. Miner
Mr. Horn
Mr. Baker
Mr. Lovingood
Mr. Ryan
Mr. Rheinfurth
Mr. Dahm
Mr. Linsley
Mr. Wilson
Mr. Clark
Mr. W. W. Vaughan
Mr. O. H. Vaughan
Mr. R. E. Smith (20)
Mr. O. E. Smith (2)
Mr. Scoggins
Mr. Daniels
Mr. Kaufman
Mr. Dalton
Dr. Heybey
Mr. Murphree
Dr. Sperling

R-ASTR

Dr. Haeussermann (2)
Mr. Hoberg

R-LVO

Dr. Gruene

I-DIR

Mr. R. B. Young

NASA Headquarters

Office of Space Sciences & Application
Dr. Homer Newell
Dr. Morris Tepper (2)
Mr. William Spreen
Mr. M. Dubin (4)

Office of Advanced Research & Tech.

Dr. R. L. Bisplinghoff
Mr. Hildred Chesley
Dr. H. Kurzweg
Mr. Milton Ames

Office of Tracking & Data Acquisition

Mr. E. Buckley

MS-T

Mr. Roy Bland

MS-IP

MS-IPL (8)

CC-P

MS-H